

Westcoast Pectus Carinatum (WPC) Compressor



WPC Compressor - *Standard Brace*



WPC Compressor II - *Brace is used for deformity superior to the nipple line*



WPC Compressor with extension - *brace for females with significant breast development*

WESTCOAST BRACE & LIMB
5311 E. Fletcher Avenue
Tampa, FL 33617



WPC Compressor™



**Treatment for
Pectus Carinatum**

What is Pectus Carinatum?

Pectus Carinatum, which is referred to as “pigeon chest,” is one of the most common chest wall deformities. The condition causes the sternum to protrude, with a narrow depression along the sides of the chest. This gives the chest a bowed-out appearance similar to that of a pigeon. People with pectus carinatum generally develop normal hearts and lungs, but the deformity may prevent these from functioning optimally. There is evidence that pectus carinatum may prevent complete expiration of air from the lungs in children. These young people may have a decrease in stamina, even if they do not recognize it.



The condition seems to appear more often in boys rather than girls. Although it appears more frequently in boys, it typically develops later for them than it does for girls.

Apart from the possible physiological consequences, pectus deformities can have a significant psychological impact. Some children live happily with pectus carinatum. For others, the shape of the chest can damage their self-image and self-confidence.

Treatment

There are two forms of treatment of Pectus Carinatum – surgery or external bracing. In severe cases, surgery may be used to restore normal chest contour. In moderate cases, a brace such as the WPC Compressor™ can assist with correcting the condition.

What is the right method of choice; Surgical vs. Non-Surgical Bracing.

It all begins with an evaluation by the child's physician, who then will refer the patient to a surgeon, who has experience in the management of congenital chest wall malformations. Together as a team you will decide which treatment option is best. Following the evaluation, the physician will then provide a diagnosis and a prescription for an evaluation by an Orthotist (bracing professional).

Bracing

When and how long will the child need to wear a brace? Bracing is another option for treating Pectus Carinatum. The patient will be required to wear it during the day, during physical activity as well as to bed. The brace will be effective only if it is worn at all times. The more the child wears the brace, the greater effect it will have. The length of wearing the brace may vary, however the average is for about a year. The results of the bracing treatment varies and is also dependent on the compliance of wearing the brace.

Before and After



For additional information about the WPC Compressor, please contact Greg Bauer, CPO or Brian Whitacre at Westcoast Brace & Limb at 1(888)552-2555 or visit our website at www.wcbl.com. **For practitioners, please refer to our website for measuring forms and the L Codes.**